In vaginal candidosis:

TRADEMARK

the elegant way
to treat an inelegant problem ketoconazole



2 oral tablets once daily for 5 days is all it takes today to effectively cure the problem

Remember:

A.

oral convenience improves patient compliance and so reduces the chance of relapse.

Presentation: white, flat, half-scored uncoated tablets marked. Janssen: on one side and K / 200 on the reverse. Each tablet contains 200 mg ketoconazole. Dosage (for vaginal candidosis only): two tablets (400 mg) once daily for 5 days. For maximal absorption Nizoral should be taken with meals. Nizoral is contra-indicated in pregnancy. Precautions: the use of agents which reduce gastric acidity (anti-cholinergic drugs, antacids, H₂ blockers) should be avoided and, if indicated, such drugs should be taken not less than 2 hours after Nizoral etcoconazole, when given together with cyclosporin A results in increased blood levels of cyclosporin A. It is important that blood levels of cyclosporin A are monitored if the two drugs are given concomitantly. Side-effects: nausea, skin rash, headache and pruritus may occasionally be observed. Alterations in liver function tests have occurred in patients on ketoconazole; these changes may be transient. Cases of hepatitis have been reported with an incidence of about 1 per 10,000 patients. Some of these may represent an idiosyncratic adverse reaction to the drug. This should be borne in mind in patients on long-term therapy. If a patient develops jaundice or any symptoms suggestive of hepatitis. treatment with ketoconazole should be stopped. Mild asymptomatic increases of liver enzyme levels, on the other hand, do not necessitate discontinuation of the treatment.

References: Tooley, et al.: The Practitioner **229**, 655 (1985) Benussi, et al.: Curr. Ther. Res. **31**(4), 511 (1982)



PHARMACEUTICA

B-2340 Beerse, Belgium


world leader in antimycotic research

ABC OF HYPERTENSION

A short guide to a complex and rapidly developing topic, the second, revised edition of ABC of Hypertension reviews up to date information from all recent clinical trials on hypertension and includes the latest recommendations of the British Hypertension Society. With its comprehensive survey of drug treatment – including advice on treating hypertension in pregnancy and in the elderly – this is an invaluable source of facts and guidance for practising doctors and medical students.

Chapters include

- Blood pressure measurement in infancy and childhood
- Drug treatment
- Blood pressure reduction in the elderly
- Hypertension in pregnancy
- Detecting hypertensive patients
- Management of hypertension in general practice



Price: Inland £4.95; Abroad £6.50; USA\$10.50 BMA members: Inland £4.45; Abroad £6.00; USA\$9.50 including postage, by air abroad

Payment must be enclosed with order



ON TREES

In the prehistoric days of medicine some drugs could be found growing on trees. Today that's a bit different.

The therapeutic drugs of our times are being unmasked from nature rather more scientifically.

By thorough bio-medical research into the fundamental processes of life. By studying exactly why and how these processes sometimes go wrong.

And so nature shows man where and how he can intervene and defend himself.
With precision drugs that will selectively and safely correct what went wrong.
Today's drugs grow from man's scientific knowledge of nature — but no longer on trees.